


The next two questions pertain to the situation described below.

A metal ring, in the page, is in a region of uniform magnetic field pointing out of the page as shown in the figure below.



- 1) If the ring moves to the right (in the direction shown by the arrow) at a constant speed, what is the direction of the induced current in the ring?
 - a. No current is induced.
 - b. Clockwise
 - c. Counterclockwise


- 2) If we now decrease the magnetic field at a constant rate, what is the direction of the induced current in the loop?
 - a. Counterclockwise
 - b. No current is induced
 - c. Clockwise

The next two questions pertain to the situation described below.

On the horizontal plane is a pair of parallel, conducting wires separated by a distance L . The left ends are connected to a resistor R as shown in the figure.

Sliding frictionlessly on the wires is a conducting bar. The resistances of the wires and bar are negligible.

A uniform magnetic field B of magnitude B is applied perpendicular to the page.



3) The conducting bar is pulled to the right at a constant speed $v = 12 \text{ m/s}$.

As the bar moves, a constant current $I = 0.2 \text{ A}$ flows in the direction indicated by the arrow.

What is the magnitude B of the magnetic field?



- a. $B = 3.2 \text{ T}$
- b. $B = 0.66 \text{ T}$
- c. $B = 0.47 \text{ T}$
- d. $B = 0.94 \text{ T}$
- e. $B = 0.31 \text{ T}$

4) The direction of the magnetic field

- a. is into the page.
- b. is out of the page.
- c. cannot be determined.

5) A ring is placed on the page and in a uniform magnetic field of strength $B = 1 \text{ T}$ pointing out of the page (see figure).

At time zero the ring has a radius $r_1 = 2 \text{ cm}$. The ring expands at a constant rate for 10 seconds until it reaches a radius of $r_2 = 12 \text{ cm}$.



This is a bad question not to have been asked.

What is flux through the loop at $t = 0 \text{ s}$ and $t = 10 \text{ s}$?


- a. $\Phi_1 = 0.00126 \text{ T} \cdot \text{m}^2$ and $\Phi_2 = 0.0452 \text{ T} \cdot \text{m}^2$
- b. $\Phi_1 = 4 \times 10^{-4} \text{ T} \cdot \text{m}^2$ and $\Phi_2 = 0.0144 \text{ T} \cdot \text{m}^2$
- c. $\Phi_1 = 0.126 \text{ T} \cdot \text{m}^2$ and $\Phi_2 = 0.754 \text{ T} \cdot \text{m}^2$

The next three questions pertain to the situation described below.

Consider the electromagnetic wave shown in the diagram.

The wave is propagating through a vacuum along the z -axis.

The electric field of the wave is parallel to the y -axis. It is a sine wave of amplitude 15 V/m. The accompanying magnetic field is parallel to the x -axis. See Figure (which is a snapshot at a particular instant).



6) What is the frequency f of this electromagnetic wave?

- a. $f = 0.138$ GHz
- b. $f = 1.304$ GHz
- c. $f = 0.652$ GHz
- d. $f = 0.435$ GHz
- e. $f = 0.326$ GHz

7) How much energy W goes through a square of area 5 m^2 perpendicular to the z -axis (parallel to the xy -plane) in one second?


- a. $W = 1.49$ J
- b. $W = 0.2$ J
- c. $W = 2.11$ J
- d. $W = 2.99$ J
- e. $W = 0.1$ J

8) This electromagnetic wave enters a medium with the index of refraction $n = 2.55$ for this frequency. What is the wavelength, λ_n , of the wave in this medium?

- a. $\lambda_n = 234.6$ cm
- b. $\lambda_n = 117.3$ cm
- c. $\lambda_n = 4.5$ cm
- d. $\lambda_n = 18$ cm
- e. $\lambda_n = 9$ cm

The next two questions pertain to the situation described below.


A plane electromagnetic wave with electric field amplitude ($E_{max} = 5.5 \text{ V/m}$) is incident on a polarizer as depicted in the figure. The **plane of electric field** indicates the plane in which the electric field lies. It makes an angle of 55 degrees with the transmission axis of the polarizer at P . The whole system is in a vacuum.



- 9) What is the amplitude of the electric field E_P immediately after passing through the first linear polarizer at P ?
- a. $E_P = 1.8 \text{ V/m}$
 - b. $E_P = 3.7 \text{ V/m}$
 - c. $E_P = 10 \text{ V/m}$
 - d. $E_P = 3.2 \text{ V/m}$
 - e. $E_P = 4.5 \text{ V/m}$
- 10) What is the intensity I of the light beyond the second polarizer at Q in terms of the intensity I_0 of the incident light?
- a. $I = 0.57I_0$
 - b. $I = 0.33I_0$
 - c. $I = 0.82I_0$
 - d. $I = 0.19I_0$
 - e. $I = 0.11I_0$

11) A person is standing a distance $D = 7$ m in front of a flat, vertical mirror. The distance from the ground to his eyes is $H = 1.9$ m.


An object is placed on the ground a distance $d = D/2 = 3.5$ m in front of the mirror.



At what height h should the **bottom** of the mirror be so that the person can see the bottom of the object?

- a. $h = 3.68$ m
- b. $h = 0.633$ m
- c. $h = 0.317$ m


12) A candle is placed in front of a convex mirror with radius R as shown in the figure.



Which ray trace is **incorrect** for a ~~principle~~ principal ray?

- a. Ray 1
- b. Ray 2
- c. Ray 3

- 13) A candle, with height $h_o = 10$ cm, is placed $d_o = 60$ cm in front of a concave mirror with radius of curvature $R = 30$ cm, as shown.




What is the magnification, m , of the image in the **concave** mirror?

- a. $m = -1$
- b. $m = -0.2$
- c. $m = -0.33$

- 14) A real object sits 6 cm in front of a mirror.

The object height is $h_o = 3$ cm.

An upright image is produced $|d_i| = 18$ cm away from the mirror.




What is the mirror?

- a. A convex mirror with the focal length (absolute value) 12 cm
- b. A convex mirror with the focal length (absolute value) 4.5 cm.
- c. A concave mirror with the focal length (absolute value) 12 cm
- d. A concave mirror with the focal length (absolute value) 4.5 cm
- e. A concave mirror with the focal length (absolute value) 9 cm

The next two questions pertain to the situation described below.

Consider a glass prism in the shape of a right triangle that makes an angle $\alpha = 66^\circ$, as shown.

The glass has index of refraction $n_{red} = 1.5$ and $n_{blue} = 1.53$ for red and blue light, respectively.



15) A ray of red, monochromatic light travelling in air to the right hits the surface of the prism at 90° , as shown in the figure. What is the angle θ_r at which the light emerges?


- a. $\theta = 54.78^\circ$
- b. $\theta = 42.87^\circ$
- c. $\theta = 24^\circ$
- d. $\theta = 37.6^\circ$
- e. $\theta = 66^\circ$

16) Now, a ray of white light hits the surface of the prism at 90° . In what order, from top to bottom do the different colored rays emerge?

- a. Blue ray at the top, red ray at the bottom
- b. Red and blue rays at the same angle
- c. Red ray at the top, blue ray at the bottom

The next three questions pertain to the situation described below.

A converging lens of focal length $f = 12.5$ cm, in air ($n = 1$), is placed at the center of a fish tank $\ell = 122$ cm long. The right-hand end of the fish tank is painted to make a screen.




17) Where should a light bulb be placed to produce a focused, real image on the screen inside the fish tank?

- a. $d_o = 12.5$ cm
- b. $d_o = 15.72$ cm
- c. $d_o = 10.37$ cm
- d. $d_o = 1.26$ cm
- e. $d_o = 0.83$ cm

18) The image on the screen is

- a. none of these.
- b. inverted.
- c. upright.

19) The fish tank is filled with water.




To obtain a focused image of the light bulb on the screen you must

- increase the distance between the light bulb and the lens.
- decrease the distance between the light bulb and the lens.
- leave the light bulb in the same location as when the fish tank was empty.

The next two questions pertain to the situation described below.

Consider the following candle-lens case:

A candle is placed $d_o = 11$ cm to the left of a diverging lens of focal length $f_2 = -5.5$ cm as shown.



20) Which of the following statements is true about the image formed by this lens:


- Statement A: Virtual, Upright
- Statement B: Real, Inverted
- Statement C: Virtual, inverted

- Statement B*
- Statement C*
- Statement A*


21) The image distance is

- $d_i = -11$ cm
- $d_i = -3.67$ cm
- $d_i = -16.5$ cm
- $d_i = 3.67$ cm
- $d_i = 11$ cm

22) Initially, there is a converging lens alone. The real image of an object placed 4 cm in front of it makes a real image 20 cm behind this convex lens.




Now, a converging lens of focal length 6 cm is placed 16 cm from the lens as shown below.



The final image produced by this two-lens system is located:

- a. 2.4 cm to the right of the second lens.
- b. 2.4 cm to the left of the second lens.
- c. 12 cm to the left of the second lens.
- d. 6 cm to the right of the second lens.
- e. 12 cm to the right of the second lens.

23) Susan has difficulty seeing distant objects. She requires corrective contact lenses.



Susan's far-point is $d_{far} = 45$ cm.

What should her corrective lens prescription be to see a koala very far away?

Remember: a diopter is $P = 1/f$ where f is measured in meters.

- a. 2.2 diopters
- b. -0.45 diopters
- c. 0.45 diopters
- d. -4.4 diopters
- e. -2.2 diopters